

# ON BRANCHED COVERING REPRESENTATION OF 4-MANIFOLDS

RICCARDO PIERGALLINI

*Scuola di Scienze e Tecnologie  
Università di Camerino – Italy*

riccardo.piergallini@unicam.it

DANIELE ZUDDAS

*Korea Institute for Advanced Study  
Seoul – Republic of Korea*

zuddas@kias.re.kr

## Abstract

We provide new branched covering representations for bounded and/or non-compact 4-manifolds, which extend the known ones for closed 4-manifolds.

Assuming  $M$  to be a connected oriented PL 4-manifold, our main results are the following: (1) if  $M$  is compact with (possibly empty) boundary, there exists a simple branched cover  $p: M \rightarrow S^4 - \text{Int}(B_1^4 \cup \dots \cup B_n^4)$ , where the  $B_i^4$ 's are disjoint PL 4-balls,  $n \geq 0$  is the number of boundary components of  $M$ ; (2) if  $M$  is open, there exists a simple branched cover  $p: M \rightarrow S^4 - \text{End } M$ , where  $\text{End } M$  is the end space of  $M$  tamely embedded in  $S^4$ .

In both cases, the degree  $d(p)$  and the branching set  $B_p$  of  $p$  can be assumed to satisfy one of these conditions: (1)  $d(p) = 4$  and  $B_p$  is a properly self-transversally immersed locally flat PL surface; (2)  $d(p) = 5$  and  $B_p$  is a properly embedded locally flat PL surface. In the compact (resp. open) case, by relaxing the assumption on the degree we can have  $B^4$  (resp.  $R^4$ ) as the base of the covering.

We also define the notion of branched covering between topological manifolds, which extends the usual one in the PL category. In this setting, as an interesting consequence of the above results, we prove that any closed oriented *topological* 4-manifold is a 4-fold branched covering of  $S^4$ . According to almost-smoothability of 4-manifolds, this branched cover could be wild at a single point.

*Keywords:* branched coverings, wild branched coverings, 4-manifolds.

*AMS Classification:* 57M12, 57M30, 57N13.

## Introduction

In [11], Montesinos proved that any oriented 4-dimensional 2-handlebody is a 3-fold simple cover of  $B^4$  branched over a ribbon surface. In [17], based on this result and on covering moves for 3-manifolds (see [16]), the first author proved that every closed connected oriented PL 4-manifold  $M$  is a four-fold simple cover of  $S^4$  branched over an immersed locally flat PL surface, possibly having a finite number of transversal double points. Then, in [8] the double points of the branch set were shown to be removable after stabilizing the covering with an extra fifth sheet, in order to get an embedded locally flat PL surface. This partially solves Problem

4.113 (A) of Kirby's list [9], but it is still unknown whether double points of the branch set can be removed without stabilization.

It is then natural to ask whether such results can be generalized to arbitrary compact 4-manifolds with (possibly disconnected) boundary and to open 4-manifolds. Moreover, it is intriguing to explore what we can do in the TOP category, namely for compact topological 4-manifolds.

The aim of the present article is to answer these questions. In the PL category, we prove Theorems 1.4 and 1.5 below in the compact case, as well as Theorems 1.6 and 1.8 in the open case. Then, by compactifying coverings, we obtain Theorem 2.3, which provides a similar representation result for *topological* 4-manifold in terms of (possibly wild) topological branched coverings, according to Definitions 2.1 and 2.2.

These results were inspired by Guido Pollini's PhD thesis [18], written under the advise of the first author. We are grateful to Guido for his contribution.

We will always adopt the PL point of view if not differently stated, referring to [19] for the basic definitions and facts concerning PL topology. However, all our results in the PL category also have a smooth counterpart, being  $\text{PL} = \text{DIFF}$  in dimension four.

## 1. Definitions and results in the PL category

We recall that a *branched covering*  $M \rightarrow N$  between compact PL manifolds is defined as a non-degenerate PL map that restricts to a (finite degree) ordinary covering over the complement of a codimension two closed subpolyhedron of  $N$ . This is the usual specialization to compact PL manifolds of the very general topological notion of branched covering introduced by Fox in his celebrated paper [5] (see also [14]).

First of all, we extend the above definition to non-compact PL manifolds. In doing so, we also remove the finiteness assumption on the degree. This will be useful in Theorem 1.8, where we need infinitely many sheets.

**DEFINITION 1.1.** We call a non-degenerate PL map  $p: M \rightarrow N$  between PL  $m$ -manifolds with (possibly empty) boundary a  $d$ -fold *branched covering*, provided the following two properties are satisfied: (1) every  $y \in N$  has a compact connected neighborhood  $C \subset N$  such that all the connected components of  $p^{-1}(C)$  are compact; (2) the restriction  $p|_1: M - p^{-1}(B_p) \rightarrow N - B_p$  over the complement of an  $(m-2)$ -dimensional closed subpolyhedron  $B_p \subset N$  is an ordinary covering of degree  $d \leq \infty$ .

More precisely, by  $B_p$  we denote the minimal subpolyhedron of  $N$  satisfying property (2). This is unique and is called the *branch set* of the branched covering  $p$ . The degree  $d = d(p)$  coincides with the maximum cardinality of the fibers  $p^{-1}(y)$  with  $y \in N$  and it is called the *degree* of the branched covering  $p$ . In fact, when  $d(p)$  is finite, then  $y \in B_p$  if and only if  $p^{-1}(y)$  has cardinality less than  $d(p)$ .

We remark that property (1) in the above definition implies (and, in our situation, it is equivalent to) the completeness of  $p$  in the sense of Fox [5], therefore  $p$  is the Fox completion (cf. [5] or [14]) of its restriction  $p|_1: M - p^{-1}(B_p) \rightarrow N - B_p$ . As such,  $p$  is completely determined, up to PL homeomorphisms, by the inclusion  $B_p \subset N$  and by the ordinary covering  $p|_1: M - p^{-1}(B_p) \rightarrow N - B_p$ , or equivalently, by the associated *monodromy* homomorphism  $\omega_p: \pi_1(N - B_p) \rightarrow \Sigma_{d(p)}$ . Finally,  $p$  is called

a *simple* branched covering if the monodromy  $\omega_p(\mu)$  of any meridian  $\mu \in \pi_1(N - B_p)$  around  $B_p$  is a transposition (in general, it decomposes into disjoint cycles of finite order). In the special case when  $N$  is simply connected, the group  $\pi_1(N - B_p)$  is generated by such meridians and the monodromy can be encoded by labeling (a diagram of)  $B_p$  with the transpositions corresponding to them.

According to the above definitions and notations, we collect the results mentioned in the introduction in the following statement.

**THEOREM 1.2** ([17, 8]). *Every closed connected oriented PL 4-manifold  $M$  can be represented by a simple branched covering  $p: M \rightarrow S^4$ , with degree  $d(p)$  and branch set  $B_p \subset S^4$  satisfying one of the following conditions:*

- (a)  $d(p) = 4$  and  $B_p$  is a self-transversally immersed locally flat PL surface;
- (b)  $d(p) = 5$  and  $B_p$  is an embedded locally flat PL surface.

Next theorems represent the extensions of the previous one to bounded and open 4-manifolds, respectively, that we will prove in this paper. In order to state and prove them, we recall the notion of ribbon surface in  $B^4$  and introduce the ribbon extendability property for simple branched coverings of  $S^3$ .

A properly embedded PL surface  $S \subset B^4$  is a *ribbon surface* if and only if it can be realized by pushing inside  $B^4$  the interior of a PL immersed surface  $S' \subset S^3 = \partial B^4$ , whose only self-intersections consist of transversal double arcs like the one depicted in Figure 1. Up to PL isotopy of ribbon surfaces in  $B^4$  the surface  $S$  is uniquely determined by the surface  $S'$ , which is called the 3-dimensional diagram of  $S$ , and in the Figures we will always draw the latter to represent the former.




FIGURE 1. A self-intersection arc in the diagram of a ribbon surface.

**DEFINITION 1.3.** A simple branched covering  $p: M \rightarrow S^3$  is defined to be *ribbon extendable* if it can be extended to a simple branched covering  $q: W \rightarrow B^4$  whose branch set  $B_q \subset B^4$  is a ribbon surface (which immediately implies that  $M = \partial W$ ,  $B_p = \partial B_q \subset S^3$  is a link, and  $d(p) = d(q)$ ). For the sake of convenience, we also call ribbon extendable any simple branched cover  $p: M \rightarrow S_1^3 \cup \dots \cup S_n^3$  that is a disjoint union of ribbon extendable coverings.

We observe that the above definition is invariant under equivalence of  $p$  up to PL homeomorphisms. Hence, ribbon extendability of  $p: M \rightarrow S^3$  can be expressed in terms of the labeled branch set  $B_p$  by requiring that it is a labeled link in  $S^3$  bounding a labeled ribbon surface in  $B^4$ .

When using simple branched coverings of  $S^3$  to represent closed connected oriented 3-manifolds, ribbon extendability arises quite naturally and it is not so restrictive. In fact, it is satisfied by any branched covering representation of such a 3-manifold derived from an integral surgery description of it by the procedure given in [11] (cf. also [4]) or by the more effective one provided in [1, 2] (see Section 3 below).

**THEOREM 1.4.** *Every compact connected oriented PL 4-manifold  $M$  with  $n$  boundary components can be represented by a simple branched covering  $p: M \rightarrow S^4 - \text{Int}(B_1^4 \cup \dots \cup B_n^4)$  satisfying property (a) or (b) as in Theorem 1.2, with the  $B_i^4$ 's pairwise disjoint standard PL 4-balls in  $S^4$  and  $B_p$  a bounded surface properly immersed or embedded in  $S^4 - \text{Int}(B_1^4 \cup \dots \cup B_n^4)$ . Moreover, the restriction of  $p$  to the boundary can be required to coincide with any given ribbon extendable simple branched covering  $b: \partial M \rightarrow \partial B_1^4 \cup \dots \cup \partial B_n^4$  with  $d(b) = d(p)$ .*

In Theorem 1.4, if the boundary is connected and non-empty, that is  $n = 1$ , we have a simple branched covering  $p: M \rightarrow B^4$ . By relaxing the constraint on the degree, we can always require that the base of the covering be  $B^4$ , even if  $M$  has more than one boundary component.

**THEOREM 1.5.** *Every compact connected oriented PL 4-manifold  $M$  with  $n \geq 2$  boundary components is a  $3n$ -fold simple covering of  $B^4$  branched over a properly embedded locally flat PL surface in  $B^4$ . Moreover, the restriction of the covering to the boundary can be required to coincide with any given  $3n$ -fold ribbon extendable simple branched covering of  $S^3$ .*

For a non-compact manifold  $M$ , we denote by  $\text{End } M$  the end space of  $M$ , that is, the inverse limit of the inclusion system of component spaces  $C(M - K)$  with  $K$  varying on the compact subspaces  $K \subset M$  (see [6]). Since  $\text{End } M$  is a compact totally disconnected metrizable space, possibly containing a Cantor set, it can be embedded in  $R$ .

**THEOREM 1.6.** *Every open connected oriented PL 4-manifold  $M$  can be represented by a simple branched covering  $p: M \rightarrow S^4 - \text{End } M$  satisfying property (a) or (b) as in Theorem 1.2, with  $\text{End } M$  embedded in  $S^4$  as a tame totally disconnected subspace (in particular, we can have  $\text{End } M \subset S^1 \subset S^4$ ) and  $B_p$  an unbounded surface properly immersed or embedded in  $S^4 - \text{End } M$ .*

In the special case when  $M$  has only one end, that is  $\text{End } M$  consists of a single point, then Theorem 1.6 tells us that  $M$  is a simple branched covering of  $R^4$ . As a direct consequence we have the following corollary.

**COROLLARY 1.7.** *For every exotic  $R_{\text{ex}}^4$  there is a simple branched covering  $p: R_{\text{ex}}^4 \rightarrow R^4$  to the standard  $R^4$ , satisfying property (a) or (b) as in Theorem 1.2.*

In the same spirit of Theorem 1.5, we have a similar result for open 4-manifolds. Namely, by relaxing the constraint on the degree as above, we can always require that the base of the covering is  $R^4$ , even if  $M$  has more than one end.

**THEOREM 1.8.** *Every open connected oriented PL 4-manifold  $M$  with more than one end is a  $3n$ -fold simple covering of  $R^4$  branched over a properly embedded locally flat PL surface in  $R^4$ , with  $n = \min\{\aleph_0, |\text{End } M|\}$ .*

The theorems above can be combined in various ways, by including in a single statement different points of view. In particular, we limit ourselves to consider next Theorems 1.9 and 1.10 below. The former includes Theorems 1.4 and 1.6, while the latter includes Theorems 1.4 and 1.5, as well as Lemma 3.6 stated in Section 3. The proofs of these new theorems are nothing else than combinations of the proofs of the constituent ones, so we leave them to the reader.

**THEOREM 1.9.** *For every connected oriented PL 4-manifold  $M$  with (possibly empty) compact boundary, there exists a simple branched cover  $p: M \rightarrow S^4 - \text{Int}(B_1^4 \cup \dots \cup B_n^4) - \text{End } M$  satisfying property (a) or (b) as in Theorem 1.2, where the  $B_i^4$ 's are pairwise disjoint locally flat PL 4-balls in  $S^4$ ,  $n \geq 0$  is the number of boundary components of  $M$ , and  $\text{End } M$  is the (possibly empty) end space of  $M$  tamely embedded in  $S^4 - \text{Int}(B_1^4 \cup \dots \cup B_n^4)$ .*

**THEOREM 1.10.** *Let  $M$  be a compact connected oriented PL 4-manifold with boundary and  $b: \partial M \rightarrow S_1^3 \cup \dots \cup S_k^3$  be a  $d$ -fold ribbon extendable simple branched covering over a disjoint union of 3-spheres, with  $k \geq 1$  and  $d \geq 4$  (resp.  $d \geq 5$ ). Then  $b$  can be extended to a  $d$ -fold simple branched covering  $p: M \rightarrow S^4 - \text{Int}(B_1^4 \cup \dots \cup B_k^4)$ , whose branch set  $B_p$  is a properly self-transversally immersed (resp. a properly embedded) locally flat PL surface.*

## 2. Branched coverings in the TOP category

In order to deal with topological 4-manifolds, we need a more general notion of branched covering, not requiring PL structures and admitting a possibly wild branch set.

**DEFINITION 2.1.** We call a continuous map  $p: M \rightarrow N$  between topological  $m$ -manifolds with (possibly empty) boundary a *tame topological branched covering* if it is locally modeled on PL branched coverings, meaning that for every  $y \in N$  there exists a local chart  $V$  of  $N$  at  $y$  and pairwise disjoint local charts  $U_i$  of  $M$  at all the  $x_i \in p^{-1}(y)$ , such that  $p^{-1}(V) = U = \cup_i U_i$  and  $p|_U: U \rightarrow V$  is a PL branched covering.

The local chart  $V$  in the above definition can be replaced by an  $m$ -ball  $C$  centered at  $y$  such that  $p^{-1}(C) = \cup_i C_i$  is the union of pairwise disjoint  $m$ -balls, each  $C_i$  being centered at a point  $x_i$  of  $p^{-1}(y)$  and each restriction  $p|_{C_i}: C_i \rightarrow C$  being topologically equivalent to the cone of a PL branched covering  $S^{m-1} \rightarrow S^{m-1}$ . Using this local conical structure, one could also define the notion of topological branched covering by induction on the dimension  $m$ , starting with ordinary coverings for  $m = 1$ .

As an immediate consequence of the existence of the local models, a tame topological branched covering  $p$  is a discrete open map. Furthermore, the union of all the branch sets of the local restrictions over charts  $V$  as in the definition is an  $(m - 2)$ -dimensional (locally tame) subspace  $B_p \subset N$ , which we call the *branch set* of  $p$ , and the restriction  $p|_{M - p^{-1}(B_p)}: M - p^{-1}(B_p) \rightarrow N - B_p$  over the complement of  $B_p$  is an ordinary covering of degree  $d(p) \leq \infty$ , which we call the *degree* of  $p$ . So,  $p$  satisfies property (2) as in Definition 1.1, but with  $B_p$  being a polyhedron only locally.

On the other hand,  $p$  turns out to be complete, satisfying the condition (1) as in Definition 1.1, hence it is the Fox completion of  $p|_{M - p^{-1}(B_p)}: M - p^{-1}(B_p) \rightarrow N - B_p$  (cf. [5] or [14]). Thus, like in the PL case,  $p$  is completely determined, up to homeomorphisms, by the inclusion  $B_p \subset N$  and by the *monodromy* homomorphism  $\omega_p: \pi_1(N - B_p) \rightarrow \Sigma_{d(p)}$ . Moreover, it still makes sense to speak of meridians around  $B_p$  and to call  $p$  *simple* if the monodromy of each meridian is a transposition.

DEFINITION 2.2. We call a continuous map  $q: M \rightarrow N$  between topological  $m$ -manifolds with (possibly empty) boundary a *wild topological branched covering* if it is discrete and open,  $q^{-1}(\partial N) = \partial M$ , and the following two conditions hold: (1) every  $y \in N$  has a compact connected neighborhood  $C \subset N$  such that all the connected components of  $q^{-1}(C)$  are compact; (2) the restriction  $p = q|_M: M - q^{-1}(W_q) \rightarrow N - W_q$  over the complement of a closed nowhere dense subspace  $W_q \subset N$  is a tame topological branched covering.

We always assume  $W_q$  to be minimal with the property required in the above definition, and call it the *wild set* of  $q$ . Of course  $q$  is actually wild only if  $W_q \neq \emptyset$ , otherwise it is a tame topological branched covering.

For a wild topological branched covering  $q: M \rightarrow N$ , with  $p$  its tame restriction as in the definition, we call  $B_q = W_q \cup B_p$  the *branch set* of  $q$  and  $d(q) = d(p)$  the *degree* of  $q$ . By the minimality of  $W_q$  and  $B_p$ , we have  $B_q = q(S_q)$ , with  $S_q \subset M$  denoting the *singular set* of  $q$ , that is, the set of points of  $M$  where  $q$  is not a local homeomorphism. Then, Theorem 2 of [3] applies to give  $\dim S_q = \dim B_q \leq m - 2$ , which easily implies that  $\dim q^{-1}(B_q) \leq m - 2$  as well. Therefore,  $N - B_q$  and  $M - q^{-1}(B_q)$  are dense and locally connected in  $N$  and  $M$ , respectively, and so we can conclude that  $q$  is the Fox completion of the restriction  $q|_M: M - q^{-1}(B_q) \rightarrow N$ . Since  $q|_M: M - q^{-1}(B_q) \rightarrow N - B_q$  is an ordinary covering,  $q$  is a branched covering in the sense of Fox [5] (for  $M$  connected) and Montesinos [14], and it is completely determined, up to topological equivalence, by the inclusion  $B_q \subset N$  and the *monodromy*  $\omega_q = \omega_p: \pi_1(N - B_q) \rightarrow \Sigma_{d(q)}$ .

In the special case when  $M$  and  $N$  are compact and  $\dim W_q = 0$ , according to Montesinos in [13, Theorem 2], the Fox *compactification theorem* [5, pag. 249] can be generalized to see that  $q$  is actually the Freudenthal *end compactification* (see [6]) of its restriction  $p$  over  $N - W_q$ . In particular,  $M$  and  $N$  are the end compactifications of  $M - q^{-1}(W_q)$  and  $N - W_q$ , respectively, hence  $q^{-1}(W_q) \cong \text{End}(M - q^{-1}(W_q))$  and  $W_q \cong \text{End}(N - W_q)$ .

In light of the above definitions and recalling that any open 4-manifold admits a PL structure (is smoothable) [7], we can state our third theorem about the branched covering representation of topological 4-manifolds.

THEOREM 2.3. *Every closed connected oriented topological 4-manifold  $M$  can be represented by a topological branched covering  $q: M \rightarrow S^4$ , which is the one-point compactification of a simple PL branched covering of  $R^4$  satisfying property (a) or (b) as in Theorem 1.2. Then, the branch set  $B_q$  is the one-point compactification of a surface in  $R^4$  and the wild set  $W_q$  consists of at most a single point.*

### 3. Proofs

Our starting point is the branched covering representation provided in [1, 2] of compact connected oriented 4-dimensional 2-handlebodies up to 2-deformations. As usual, here and in the following, we call a 2-handlebody any handlebody whose all handles have index  $\leq 2$ , and a 2-deformation any sequence of handle operations (isotopy, sliding and addition/deletion of canceling handles) not involving any handle of index  $> 2$ .

Below we briefly recall the procedure described in [1, Section 3] (see also [2, Sections 6.1 and 3.4]), for deriving from any Kirby diagram  $K$  of a connected oriented 4-dimensional 2-handlebody  $H$  a labeled ribbon surface  $S_K \subset B^4$  representing a simple 3-fold covering  $p: H \rightarrow B^4$  branched over  $S_K$ .




FIGURE 2. A Kirby diagram  $K$  and the labeled ribbon surface  $S_K$ .

Let  $K \subset R^3 \cong S^3 - \{\infty\}$  be any Kirby diagram of an oriented 4-dimensional 2-handlebody  $H \cong H^0 \cup H_1^1 \cup \dots \cup H_m^1 \cup H_1^2 \cup \dots \cup H_n^2$  with a single 0-handle  $H^0$ , 1-handles  $H_i^1$  and 2-handles  $H_j^2$ . Denote by  $B_1, \dots, B_m \subset R^3$  the disjoint disks spanned by the dotted unknots of  $K$  representing the 1-handles and by  $L_1, \dots, L_n \subset R^3$  the framed components of  $K$  representing the attaching maps of the 2-handles. Then, the labeled ribbon surface  $S_K \subset R_+^4 \cong B^4 - \{\infty\}$  can be constructed as follows (look at Figure 2, where a simple Kirby diagram  $K$  and the corresponding labeled ribbon surface  $S_K$  are drawn, respectively on the left and on the right side).

PROCEDURE 3.1 (CONSTRUCTION OF  $S_K$ ).

- 1) Choose a trivializing set of crossings in the diagram of the framed link  $L = L_1 \cup \dots \cup L_n$  (the asterisked ones in Figure 2); denote by  $L' = L'_1 \cup \dots \cup L'_n \subset R^3$  the trivial link obtained by inverting those crossings, and by  $D_1, \dots, D_n \subset R^3$  a family of disjoint disks spanned by  $L'_1, \dots, L'_n$ , respectively;
- 2) let  $A_1, \dots, A_n \subset R^3$  be a family of disjoint (possibly non-orientable) narrow closed bands, each  $A_j$  having  $L'_j$  as the core and representing half the framing that  $L'_j$  inherits from  $L_j$  (by parallel transport at the crossing changes);
- 3) let  $B'_1, \dots, B'_m \subset R^3$  be a family of disjoint disks, each  $B'_i$  being parallel to  $B_i$ ;
- 4) let  $C_1, \dots, C_\ell \subset R^3$  be a family of disjoint small disks, each  $C_k$  being placed at one of the trivializing crossings and forming with the involved bands  $A_j$  a fixed pattern of ribbon intersections inside a 3-ball thickening of it, as in Figure 2;
- 5) choose a family of disjoint narrow bands  $\alpha_1, \dots, \alpha_n \subset R^3$ , each  $\alpha_j$  connecting  $A_j$  to a fixed disk  $A_0$  disjoint from all the other disks and bands, with the only constraints that it cannot meet any disk  $D_1, \dots, D_n$ , the 3-ball spanned by any pair of parallel disks  $B_i$  and  $B'_i$ , and the 3-ball thickening of any  $C_k$ ;

- 6) choose a family of disjoint narrow bands  $\beta_1, \dots, \beta_m \subset R^3$ , each  $\beta_i$  connecting  $B'_i$  to a fixed disk  $B_0$  disjoint from all the other disks and bands, with the same constraints as above;
- 7) choose a family of disjoint narrow bands  $\gamma_1, \dots, \gamma_\ell \subset R^3$ , each  $\gamma_k$  connecting  $C_k$  to the disk  $B_0$ , with the same constraints as above;
- 8) put  $A = A_0 \cup_{j=1}^n (\alpha_j \cup A_j) \subset R^3$  and  $B = B_0 \cup_{i=1}^m (\beta_i \cup B'_i) \cup_{k=1}^\ell (\gamma_k \cup C_k) \subset R^3$ ;
- 9) then,  $S_K \subset R_+^4 \subset B^4$  is the ribbon surface whose 3-dimensional diagram is given by  $A \cup B \cup B_1 \cup \dots \cup B_m$ ; in other words,  $S_K$  is obtained by pushing the interior of the connected surfaces  $A, B, B_1, \dots, B_m$  inside the interior of  $R_+^4$ , in such a way that all the ribbon intersections (formed by  $A$  passing through  $B \cup B_1 \cup \dots \cup B_m$ ) disappear;
- 10) finally, the labeling of  $S_K$  giving the monodromy of the simple 3-fold branched covering  $p: H \rightarrow B^4$  is the one determined by assigning the transpositions (1 2) and (2 3) to the standard meridians of  $A_0$  and  $B_0 \cup B_1 \cup \dots \cup B_m$ , respectively, in the 3-dimensional diagram of  $S_K$ .

The construction above depends on various choices, the significant ones being in steps 1, 5, 6 and 7. However, the labeled ribbon surfaces obtained from different choices become equivalent up to labeled isotopy of ribbon surfaces in  $B^4$  (called 1-isotopy in [1, 2]) and the covering moves  $R_1$  and  $R_2$  depicted in Figure 3, after adding to them a separate trivial disk with label (3 4). We recall that the addition of such disk represents the stabilization of the branched covering  $p$  with an extra trivial fourth sheet to give a simple 4-fold branched covering  $\tilde{p}: H \cong H \#_{\partial} B^4 \rightarrow B^4$ .

The labels  $a, b, c$  and  $d$  in Figure 3, as well as in Figures 4 to 7, are assumed to be pairwise distinct.




FIGURE 3. Covering moves for labeled ribbon surfaces.

The covering space  $H$  of any simple branched covering  $p: H \rightarrow B^4$  described by a labeled ribbon surface  $S \subset B^4$  is a 4-dimensional 2-handlebody whose handle structure is uniquely determined, up to 2-deformations, by the ribbon structure of  $S$ . Moreover, the following equivalence theorem holds (Theorem 1 in [1], Theorem 6.1.5 in [2]).

**THEOREM 3.2** ([1, 2]). *Let  $S$  and  $S'$  be two labeled ribbon surfaces in  $B^4$  representing compact connected oriented 4-dimensional 2-handlebodies as simple branched coverings of  $B^4$  of the same degree  $\geq 4$ . Then,  $S$  and  $S'$  are related by labeled isotopy of ribbon surfaces and the moves  $R_1$  and  $R_2$  in Figure 3 if and only if the handlebodies they represent are equivalent up to 2-deformations.*

For the purposes of this paper, we need to consider the implication of the above theorem on the boundary. This implication is stated in a precise way in the next

theorem, which is a restatement of Theorem 2 in [1], or Theorem 6.1.8 in [2]. In fact, handle trading and blow-up moves (see Figure 4), introduced therein in order to interpret the Kirby calculus for 3-manifolds in terms of labeled ribbon surfaces, reduce to isotopy when restricted to the boundary.




FIGURE 4. Handle trading and blow-up moves for labeled ribbon surfaces.

**THEOREM 3.3** ([1, 2]). *Let  $L$  and  $L'$  be two labeled links in  $S^3$  representing closed connected oriented 3-manifolds as ribbon extendable simple branched coverings of  $S^3$  of the same degree  $\geq 4$ . Then,  $L$  and  $L'$  are related by labeled isotopy and the moves  $B_1$  and  $B_2$  in Figure 5 if and only if the oriented 3-manifolds they represent are PL homeomorphic.*




FIGURE 5. Covering moves for labeled links bounding labeled ribbon surfaces.

Now, before proceeding with the proofs of the theorems stated in Section 1, let us prove two lemmas.

**LEMMA 3.4.** *Let  $W \cong M \times [0, 1] \cup H_1^1 \cup \dots \cup H_m^1 \cup H_1^2 \cup \dots \cup H_n^2$  be an oriented 4-dimensional 2-cobordism between closed connected oriented 3-manifolds  $M_0 = M \times \{0\}$  and  $M_1 = \partial W - M_0$ . Then, any  $d$ -fold simple branched covering  $p_0: M_0 \rightarrow S^3 \times \{0\}$  of degree  $d \geq 3$  extends to a  $d$ -fold simple branched covering  $p: W \rightarrow S^3 \times [0, 1]$ , such that  $B_p \subset S^3 \times [0, 1]$  is a properly embedded locally flat PL surface. Moreover, if  $p_0$  is ribbon extendable, then we can choose  $p$  in such a way that also  $p_1 = p|_{M_1}: M_1 \rightarrow S^3 \times \{1\}$  is ribbon extendable.*

*Proof.* This immediately follows from the main result in [4] and its proof.  $\square$

**REMARK 3.5.** If one is only interested in the existence of a 3-fold covering  $p: W \rightarrow S^3 \times [0, 1]$  as in the lemma above, without insisting that it restricts to a given covering  $p_0$ , then the following argument provides a more explicit construction.

Let  $K_0$  be a Kirby diagram representing a 4-dimensional 2-handlebody  $W_0 = H^0 \cup H_{n+1}^2 \cup \dots \cup H_\ell^2$  such that  $\partial W_0 \cong M$ . By identifying a collar  $C$  of  $\partial W_0$  in  $W_0$  with  $M \times [0, 1] \subset W$ , in such a way that  $\partial W_0$  corresponds to  $M \times \{1\}$ , we get a 4-dimensional 2-handlebody  $W_1 = W_0 \cup_{C \cong M \times [0, 1]} W = H^0 \cup H_1^1 \cup \dots \cup H_m^1 \cup H_1^2 \cup \dots \cup$

$H_n^2 \cup H_{n+1}^2 \cup \dots \cup H_\ell^2$ . Here, the handles have been reordered in the usual way, once the attaching maps of the handles of  $W$  are isotoped in  $\partial W_0$  out of the 2-handles of  $W_0$ . So, we have a Kirby diagram  $K_1$  of  $W_1$  that contains  $K_0$  as a framed sublink.

Procedure 3.1 determines a labeled ribbon surface  $S_{K_1}$ . By pushing the part of  $S_{K_1}$  corresponding to  $K_0$  a little bit more inside the interior of  $B^4$  than the rest of  $S_{K_1}$ , we can assume that for some  $r < 1$  the intersection of  $S_{K_1}$  with the 4-ball  $B_r^4 \subset \text{Int } B^4$  of radius  $r$  is a copy of  $S_{K_0}$  in  $B_r^4$ . Then, the branched covering  $q_1: W_1 \rightarrow B^4$  represented by  $S_{K_1}$  restricts to two branched coverings  $q_0: W_0 \rightarrow B_r^4$  and  $q: W \rightarrow B^4 - \text{Int } B_r^4$ . At this point, the desired 3-fold simple branched covering  $p: W \rightarrow S^3 \times [0, 1]$  is just the composition of  $q$  with the canonical identification  $B^4 - \text{Int } B_r^4 \cong S^3 \times [0, 1]$ .

**LEMMA 3.6.** *Let  $M$  be a closed connected oriented 3-manifold and assume  $d \geq 4$ . For any two  $d$ -fold ribbon extendable simple branched coverings  $p_0, p_1: M \rightarrow S^3$ , there is a  $d$ -fold simple branched covering  $p: M \times [0, 1] \rightarrow S^3 \times [0, 1]$  satisfying the following properties: 1) the restriction  $p|_{M \times \{0\}}: M \times \{0\} \rightarrow S^3 \times \{0\}$  coincides with  $p_0 \times \{0\}$ ; 2) the restriction  $p|_{M \times \{1\}}: M \times \{1\} \rightarrow S^3 \times \{1\}$  is equivalent to  $p_1 \times \{1\}$  up to PL homeomorphisms; 3) the branch set  $B_p \subset S^3 \times [0, 1]$  is a properly immersed locally flat PL surface, whose singularities (if any) consist of an even number of transversal double points. In addition, if  $d \geq 5$  there is such a branched covering  $p$  with  $B_p$  a properly embedded surface.*

*Proof.* By Theorem 3.3, the labeled links  $L_0$  and  $L_1$  representing the coverings  $p_0$  and  $p_1$ , respectively, are related by labeled isotopy and moves  $B_1$  and  $B_2$  depicted in Figure 5. Each move  $B_i$  can be realized as a composition of two iterations of the same Montesinos move  $M_i$  depicted in Figure 6, applied in opposite directions for  $i = 1$  (the two directions are equivalent for  $i = 2$ ). This is shown in Figure 7 for  $B_1$ , while it is trivial for  $B_2$  (cf. [1, page 5], or the proof of Theorem 6.2.3 in [2]).




FIGURE 6. Montesinos covering moves for labeled links.




FIGURE 7. Generating  $B_1$  move as the composition of two (opposite)  $M_1$  moves.

The labeled links  $L_0$  to  $L_1$  can be joined by a family of singular links  $L_t \subset S^3$  with  $t \in [0, 1]$ , which present a singular point at a finite (even) number of values of

$t$ , say  $t_1 < \dots < t_{2n}$ , in correspondence of the Montesinos moves, while giving an isotopic deformation of (non-singular) links in each open interval  $(t_i, t_{i+1})$  for  $i = 1, \dots, 2n-1$ . Following the argument proposed by Montesinos in [12], and then used in [17], things can be arranged in such a way that  $S = \cup_{t \in [0,1]} (L_t \times \{t\}) \subset S^3 \times [0, 1]$  is a labeled locally flat PL surface with a cusp singularity (the cone of a trefoil knot) for each move  $M_1$  and a node singularity (a transversal double point) for each move  $M_2$ . This is suggested by Figure 8.




FIGURE 8. Singularities of the branch surface deriving from Montesinos moves.

Then, the labeled surface  $S$  determines a  $d$ -fold simple branched covering  $q: M \times [0, 1] \rightarrow S^3 \times [0, 1]$ , whose restrictions over  $S^3 \times \{0\}$  and  $S^3 \times \{1\}$  are equivalent to  $p_0 \times \{0\}$  and  $p_1 \times \{1\}$ , respectively, up to PL homeomorphisms. In particular, there exists a PL homeomorphism  $h: M \rightarrow M$  such that  $q|_{M \times \{0\}} \circ (h \times \{0\}) = p_0 \times \{0\}$ , and we can replace  $q$  by  $q \circ (h \times \text{id}_{[0,1]})$  to have the restriction over  $S^3 \times \{0\}$  coinciding with  $p_0 \times \{0\}$  as required.

Here, cusp singularities come in pairs, each pair corresponding to two opposite moves  $M_1$  and hence consisting of cones of a left-handed and a right-handed trefoil knot. Then, since  $d \geq 4$ , the technique described in [17] applies in the present context as well, being essentially local in nature, in order to remove all the (pairs of) cusp singularities (see [8] for a different approach). As the result we get a new labeled surface  $S'$  representing a  $d$ -fold simple branched covering  $p: M \times [0, 1] \rightarrow S^3 \times [0, 1]$ , such that  $B_p = S'$  is a properly immersed locally flat PL surface whose singularities (if any) are transversal double points. Moreover, as shown in [8], if  $d \geq 5$  transversal double points can also be removed in pairs from  $B_p$  to give a properly embedded locally flat PL surface.  $\square$

At this point, we are ready to prove our main results.

*Proof of Theorem 1.4.* The existence of a branched covering  $b$  as in the second part of the statement is guaranteed by Procedure 3.1 applied to Kirby diagrams representing (4-dimensional 2-handlebodies bounded by) the components of  $\partial M$ . So, we can directly assume that  $b$  is given. We denote simply by  $d = 4$  or  $5$ , depending on the property (a) or (b) we desire, the degree  $d(b)$  of such covering.

Let us start with the case  $n = 1$ , when  $\partial M$  is connected. Given any handlebody decomposition  $H$  of  $M$  with a single 0-handle and no 4-handles, let  $M'$  be the union of the 0-handle and the 1-handles of  $H$  and put  $M'' = \text{Cl}(M - M')$ . By dualizing the 2-handles and the 3-handles of  $H$ , we can think of  $M''$  as an oriented 2-cobordism from  $\partial M$  to  $\partial M'$ .

Now,  $M'$  is the boundary connected sum of a certain number  $k$  of copies of  $S^1 \times B^3$ , hence it admits a standard representation as a 2-fold branched covering of  $B^4$ . This can be stabilized to a simple  $d$ -fold covering  $p': M' \rightarrow B^4$  branched over a ribbon surface (in fact, the union of  $k + d - 1$  separated trivial disks, with monodromies  $(12), \dots, (12), (23), (34)$  and possibly  $(45)$ , depending on  $d$ ). On the other hand, Lemma 3.4 allows us to extend the given branched covering  $b: \partial M \rightarrow \partial B_1^4 \cong \partial B_1^4 \times \{0\} \cong S^3 \times \{0\}$  to a  $d$ -fold simple covering  $p'': M'' \rightarrow \partial B_1^4 \times [0, 1] \cong S^3 \times [0, 1]$ , such that the restriction  $p'_1 = p''_1: \partial M' \rightarrow S^3 \times \{1\} \cong S^3$  is ribbon extendable. So, we can use Lemma 3.6 to get a  $d$ -fold simple branched covering  $q: \partial M' \times [0, 1] \rightarrow S^3 \times [0, 1]$  satisfying (a) or (b) and such that, with the obvious canonical identifications, the restriction  $q_0 = q_1: \partial M' \times \{0\} \rightarrow S^3 \times \{0\}$  coincides with  $p''_1$ , while the restriction  $q_1 = q_1: \partial M' \times \{1\} \rightarrow S^3 \times \{1\}$  is equivalent to the restriction  $p'_\partial = p'_1: \partial M' \rightarrow S^3$  up to PL homeomorphisms.

Then, we can glue together the coverings  $p'$  and  $p''$  through  $q$ , by identifying the corresponding restrictions, to obtain a  $d$ -fold simple branched covering  $p: M \rightarrow B^4$  with the property (a) or (b). In fact, according to [10], the result of the gluing is always PL equivalent to  $M \text{ rel } \partial M$ , no matter what the PL homeomorphism occurring in the identification between  $q_1$  and  $p'_\partial$  is. This concludes the proof of the case  $n = 1$ .

The case  $n > 1$  can be reduced to  $n = 1$  as follows. Denote by  $C_1, \dots, C_n$  the connected components of  $M$ . For every  $i = 1, \dots, n$ , we consider the restriction  $b_i = b_1: C_i \rightarrow \partial B_i^4$  and a  $d$ -fold simple covering  $q_i: W_i \rightarrow B_i^4$  branched over a ribbon surface  $B_{q_i} \subset B_i^4$  that extends  $b_i$ . Then, we enlarge the 4-balls  $B_1^4, \dots, B_n^4$  to disjoint PL 4-balls  $\widehat{B}_1^4, \dots, \widehat{B}_n^4 \subset S^4$  with a collar of their boundary, and each labeled surface  $B_{q_i}$  to a properly embedded labeled ribbon surface  $\widehat{B}_{q_i} \subset \widehat{B}_i^4$  by using the product structure along the collar. Let  $B^4 \subset S^4$  be a PL 4-ball obtained by attaching to  $\widehat{B}_1^4 \cup \dots \cup \widehat{B}_n^4$  an embedded 1-handle between  $\widehat{B}_i^4$  and  $\widehat{B}_{i+1}^4$  for each  $i = 1, \dots, n-1$ . These 1-handles can be chosen so that each attaching 3-ball meets  $\partial \widehat{B}_{q_1} \cup \dots \cup \partial \widehat{B}_{q_n}$  in  $d-1$  trivial arcs labeled  $(12), \dots, (d-1d)$ . Finally, we attach labeled bands running along the connecting 1-handles of  $B^4$  to get a labeled ribbon surface in  $B^4$ , as sketched in Figure 9 (where the bands labeled  $(45)$  occur only if  $d = 5$ ).




FIGURE 9. The labeled ribbon surface  $B_q$ .

This is the labeled branch set  $B_q \subset B^4$  of a  $d$ -fold simple branched covering  $q: W = W_0 \cup W_1 \cup \dots \cup W_n \rightarrow B^4$ , where  $W_0 \cong (\partial M \times [0, 1]) \cup H_1^1 \cup \dots \cup H_{n-1}^1$  is a 1-cobordism between  $\partial M = C_1 \cup \dots \cup C_n$  and  $C \cong C_1 \# \dots \# C_n$ , with the 1-handle  $H_i^1$  connecting  $C_i$  and  $C_{i+1}$ .

Since  $M$  is connected, we can assume  $W_0 \subset M$  and put  $M' = \text{Cl}(M - W_0)$ . The restriction  $q_0 = q|_{W_0}: W_0 \rightarrow B^4 - \text{Int}(B_1^4 \cup \dots \cup B_n^4)$  is a simple branched covering whose branch surface is properly embedded in  $B^4 - \text{Int}(B_1^4 \cup \dots \cup B_n^4)$ , while the restriction  $q_i: W_i \rightarrow B_i^4$  coincides with  $q_i$  by construction for every  $i = 1, \dots, n$ .

By construction, the restriction  $q_0|_{\partial M'}: \partial M' \rightarrow S^3$  is ribbon extendable, bounding the covering  $q$ . Hence, by the case  $n = 1$  proved above, we can extend such restriction to a simple covering  $p': M' \rightarrow B^4$  satisfying property (a) or (b). Then, to obtain the wanted branched covering  $p: M \rightarrow S^4 - \text{Int}(B_1^4 \cup \dots \cup B_n^4)$ , we just glue  $q_0$  and  $p'$  together by identifying their restrictions over  $S^3$ .  $\square$

*Proof of Theorem 1.6.* By a standard argument, it is possible to construct an infinite family  $\{M_i\}_{i \geq 0}$  of (non-empty) 4-dimensional compact connected PL submanifolds of  $M$ , such that  $M = \cup_{i \geq 0} M_i$  and  $M_i \subset \text{Int} M_{i+1}$  for every  $i \geq 0$ . Then, we put  $W_0 = M_0$  and  $W_i = \text{Cl}(M_i - M_{i-1})$  for every  $i \geq 1$ , and note that these are 4-dimensional compact PL submanifolds of  $M$ . Furthermore, for every  $i \geq 1$ , we can assume that each component  $C$  of  $W_i$  shares exactly one boundary component with  $M_{i-1}$  (otherwise, if there are more shared components, we connect them by attaching to  $M_{i-1}$  some 1-handles contained in  $C \cap \text{Int} M_i$ ). Let  $\{C_v\}_{v \in V}$  be the set of all components of all the  $W_i$ 's, and  $\{B_e\}_{e \in E}$  be the set of all their boundary components. We can think of  $V$  and  $E$  as the sets of vertices and edges of a graph  $T$ , respectively, with the edge  $e \in E$  joining the vertices  $v, w \in V$  if and only if  $C_v$  and  $C_w$  share the boundary component  $B_e$ . Actually, the above assumption about the intersection of the components of each  $W_i$  and the corresponding  $M_{i-1}$ , implies that  $T$  is a tree. We assume  $T$  rooted at the vertex  $v_0$  with  $C_{v_0} = W_0$  and orient the edges of  $T$  starting from  $v_0$ , so that each vertex  $v \neq v_0$  has a single incoming edge  $e_0^v$  and a non-empty set of outgoing edges  $e_1^v, \dots, e_{n(v)}^v$  (we will use this notation also for the edges outgoing from  $v_0$ ). According to this setting, the components of  $W_i$  are the  $C_v$  such that  $d(v, v_0) = i$ , where  $d$  denotes the edge distance in  $T$ . Moreover, for each such component  $C_v$  we have  $\partial C_v = B_{e_0^v} \cup B_{e_1^v} \cup \dots \cup B_{e_{n(v)}^v}$ , with  $B_{e_0^v}$  the unique boundary component shared with  $W_{i-1}$  if  $i \geq 1$ , while the boundary components  $B_{e_1^v}, \dots, B_{e_{n(v)}^v}$  are shared with  $W_{i+1}$ . In light of these facts, it is not difficult to see that  $\text{End} M \cong \text{End} T$ , with end points bijectively corresponding to infinite rays in  $T$  starting from  $v_0$ .

Now, based on the same tree  $T$ , we want to construct a similar pattern in  $S^4$  consisting of families  $\{C'_v\}_{v \in V}$  and  $\{B'_e\}_{e \in E}$ . We begin with any family  $\{B_e^4\}_{e \in E}$  of standard PL 4-balls in  $S^4$  satisfying the following properties: 1) the diameter of  $B_e^4$  vanishes when the edge distance  $d(e, v_0)$  goes to infinity; 2)  $B_{e_1^v}, \dots, B_{e_{n(v)}^v}$  are pairwise disjoint for every  $v \in V$  and contained in  $\text{Int} B_{e_0^v}^4$  if  $v \neq v_0$ . Then, we put  $C'_v = B_{e_0^v}^4 - \text{Int}(B_{e_1^v}^4 \cup \dots \cup B_{e_{n(v)}^v}^4)$  for every  $v \in V$  (assume  $B_{e_0^v}^4 = S^4$ ), and  $B'_e = \partial B_e^4$  for every  $e \in E$ . By the very definition, we have  $\partial C'_v = B'_{e_0^v} \cup B'_{e_1^v} \cup \dots \cup B'_{e_{n(v)}^v}$  for every  $v \in V$  (assume  $B'_{e_0^v} = \emptyset$ ). Moreover,  $C'_v$  and  $C'_w$  share the boundary component  $B'_e$  if and only if the edge  $e$  joins the vertices  $v, w \in V$  as above, and thus  $\text{End}(\cup_{v \in V} C'_v) \cong \text{End} T \cong \text{End} M$ .

The space  $X = S^4 - \cup_{v \in V} C'_v = \cap_{i \geq 0} \cup_{d(e, v_0) = i} B_e^4$  is tame in  $S^4$  (cf. [15]). In particular, we can have  $X \subset S^1$  by choosing each  $B_e^4$  to be a round spherical 4-ball centered at a point of  $S^1 \subset S^4$ . We can conclude that  $X \cong \text{End } M$ , being  $S^4$  the Freudenthal compactification of  $S^4 - X = \cup_{v \in V} C'_v$ .

At this point, we can define the desired branched covering  $p: M \rightarrow S^4 - X$  in three steps. First, for every  $e \in E$ , we choose a Kirby diagram  $K_e$  providing an integral surgery presentation of  $B_e$ , and denote by  $p_e: B_e \rightarrow B'_e \cong S^3$  the restriction to the boundary of the simple branched covering of  $B^4$  determined by the labeled ribbon surface  $S_{K_e}$ , stabilized to degree 4 or 5, depending on the property (a) or (b) we want to obtain for  $p$ . Then, for every  $v \in V$ , we apply Theorem 1.4 in order to extend  $p_{e_0^v} \cup p_{e_1^v} \cup \dots \cup p_{e_n^v}$ :  $\partial B_v \rightarrow \partial B'_v$  to a simple branched covering  $p_v: C_v \rightarrow C'_v$  satisfying property (a) or (b). Finally, we define  $p = \cup_{v \in V} p_v: M = \cup_{v \in V} C_v \rightarrow S^4 - X = \cup_{v \in V} C'_v$ .  $\square$

*Proof of Theorem 2.3.* Because of Theorem 1.2, it suffices to consider the case when  $M$  is not PL. Since any open 4-manifold admits a PL structure (see [7, Section 8.2]), we can apply Theorem 1.6 to the one-ended open connected oriented 4-manifold  $M - \{x\}$ , with  $x$  any point of  $M$ , in order to get a PL branched covering  $p: M - \{x\} \rightarrow R^4$  satisfying property (a) or (b). Then, the one-point compactification of  $p$  gives the wanted wild branched covering  $q: M \rightarrow S^4$ , once  $M$  and  $S^4$  are identified with the one-point compactifications of  $M - \{x\}$  and  $R^4$ , respectively.  $\square$

*Proof of Theorem 1.5.* Following the proof of Theorem 1.4 for  $n > 1$  and adopting the notations therein, we consider: the decomposition  $\partial M = C_1 \cup \dots \cup C_n$ ; the cobordism  $W_0 = (\partial M \times [0, 1]) \cup H_1^1 \cup \dots \cup H_{n-1}^1 \subset M$  between  $\partial M$  and  $\partial M' \cong C_1 \# \dots \# C_n$ , with  $M' = \text{Cl}(M - W_0)$ ; the 3-fold simple branched coverings  $q_i: W_i \rightarrow B^4$  and their restrictions to the boundary  $q_i|: C_i \rightarrow S^3$ ,  $i = 1, \dots, n$ ; the simple branched covering  $p': M' \rightarrow B^4$ .

Now, let  $q'_0: \partial M \times [0, 1] \rightarrow S^3 \times [0, 1]$  be the  $3n$ -fold simple branched covering such that  $C_i \times [0, 1]$  is given by the three sheets from  $3i - 2$  to  $3i$ , and the restriction  $q'_{0|C_i \times [0, 1]}$  coincides with  $q_i| \times [0, 1]$  up to a shifting by  $3(i - 1)$  in the numbering of the sheets. This covering  $q'_0$  can be extended to a  $3n$ -fold simple branched covering  $q''_0: W_0 \rightarrow S^3 \times [0, 1]$ , by adding separated trivial disks  $D_1, \dots, D_{n-1}$  to the labeled branch set, with  $\partial D_i \subset S^3 \times \{1\}$  and  $D_i$  labeled by  $(1 \ 3i + 1)$ . The restriction of  $q''_0$  over  $S^3 \times \{1\}$  gives a  $3n$ -fold simple branched covering  $q''_{0|}: \partial M' \rightarrow S^3 \times \{1\}$ .

Finally, we apply Theorem 1.4 (actually the strong version of Lemma 3.6 provided by it, where the restriction  $p|_{M \times \{1\}}$  coincides with the covering  $p_1 \times \{1\}$ ) to connect such restriction with the restriction over the boundary of a stabilization to degree  $3n > 5$  of the covering  $p': M' \rightarrow B^4$ . This gives the wanted simple branched covering  $p: M \rightarrow B^4$ , and concludes the proof of the first part of the statement.

For the second part, it suffices to replace the covering  $q'_0$  in the argument above by  $b \times [0, 1]$ , with  $b: \partial M \rightarrow S^3$  any given  $3n$ -fold ribbon extendable simple branched covering.  $\square$

*Proof of Theorem 1.8.* Consider the decomposition  $M = \cup_{i \geq 0} W_i$  and the families  $\{C_v\}_{v \in V}$  and  $\{B_e\}_{e \in E}$ , as in the proof of Theorem 1.6. For every  $i \geq 0$ , put  $M_i = W_i \cap W_{i+1} = \partial W_i \cap \partial W_{i+1}$  and observe that this is a closed 3-manifold with a finite number  $n_i \leq n$  of components. The sequence  $(n_i)_{i \geq 1}$  is non-decreasing, and

without loss of generality we can assume  $n_i \geq 2$  for every  $i \geq 0$ . Then, denoting by  $S_i^3$  and  $B_i^4$  respectively the 3-sphere and the 4-ball of radius  $i$  in  $R^4$ , there is a  $3n_i$ -fold simple branched covering  $b_i : M_i \rightarrow S_{i+1}^3$ , bounding a  $3n_i$ -fold simple covering of  $B_{i+1}^4$  branched over a ribbon surface. Theorem 1.10, which combines Theorems 1.4 and 1.5 proved above, gives us  $3n_i$ -fold simple coverings  $p_i : W_i \rightarrow \text{Cl}(B_{i+1}^4 - B_i^4)$  with  $i \geq 0$ , such that the restrictions of  $p_i$  over  $S_i^3$  and  $S_{i+1}^3$  respectively coincide with a  $3n_i$ -fold stabilization of  $b_{i-1}$  and with  $b_i$  (where  $b_{-1}$  is empty). At this point, we can glue the  $p_i$ 's together, up to stabilization. More precisely, we start with  $p_0$ , and then we add each  $p_i$  in order, by gluing it to the appropriate  $3n_i$ -fold stabilization of  $\cup_{j < i} p_j$ . This gives the wanted  $3n$ -fold branched covering  $p = \cup_{i \geq 0} p_i : M = \cup_{i \geq 0} W_i \rightarrow R^4$ .  $\square$

#### 4. Final remarks

We remark that all the simple branched coverings obtained in Theorems 1.4, 1.5, 1.6, 1.8 and 2.3 can be stabilized to any degree greater than the stated one. While this is obvious for branched coverings of  $S^4$  or  $B^4$  (like in Theorem 1.2), in the other cases it can be achieved by suitable covering stabilizations in the construction process.

We also remark that the arguments in the proofs of those theorems can be combined to prove various extensions of them. In particular, we have the following.

- 1) *Any non-compact connected oriented PL 4-manifold  $M$  whose boundary has only compact components, is a simple branched covering of  $S^4 - (\text{End } M \cup_{c \in C} \text{Int } B_c^4)$ , where  $\{B_c^4\}_{c \in C}$  is a family of pairwise disjoint 4-balls in  $S^4 - \text{End } M$ , indexed by the set  $C$  of the boundary components of  $M$ . The limit set of the balls  $\{B_c^4\}_{c \in C}$  is contained in  $\text{End } M \subset S^4$ .*
- 2) *Any compact connected oriented topological 4-manifold  $M$  with boundary is a simple topological branched covering of  $S^4 - \text{Int}(\cup_{c \in C} B_c^4)$  with at most one wild point.*

Finally, we observe that when  $M$  does not admit a PL structure the branch set  $B_p$  of the covering  $p : M \rightarrow S^4$  in Theorem 2.3 cannot be reduced to a locally flat PL surface properly immersed or embedded in  $S^4$ . In fact, in this case there is a single wild point in  $B_p$ , at which we concentrate all the pathological aspects of the topology of  $B_p$  and/or of the inclusion  $B_p \subset S^4$ . However, one might wonder if the situation could be simplified by diffusing the wild set  $W_p$ , or even more if such wild set could be eliminated at all, to get a tame topological branched covering at least under particular circumstances. So, we conclude with the following open problem.

**QUESTION 4.1.** *When, in representing a connected oriented topological 4-manifold  $M$  that is not PL by a simple branched covering  $p : M \rightarrow S^4$ , can we require  $B_p$  to be a topological surface wildly immersed or embedded in  $S^4$ ? Can we require  $p$  to be a tame topological branched covering, with  $B_p$  a (locally) tame 2-complex or a topological surface (locally) tamely immersed or embedded in  $S^4$ ?*

## References

- [1] I. Bobtcheva and R. Piergallini, *Covering moves and Kirby calculus*, preprint 2005, arXiv:math/0407032.
- [2] I. Bobtcheva and R. Piergallini, *On 4-dimensional 2-handlebodies and 3-manifolds*, J. Knot Theory Ramifications **21** (2012), 1250110 (230 pages).
- [3] P.T. Church, *Discrete maps on manifolds*, Michigan Math. J. **25** (1978), 351–357.
- [4] A.L. Edmonds, *Extending a branched covering over a handle*, Pacific J. of Math. **79** (1978), 363–369.
- [5] R.H. Fox, *Covering spaces with singularities*, in “Algebraic Geometry and Topology. A symposium in honour of S. Lefschetz”, Princeton University Press 1957, 243–257.
- [6] H. Freudenthal, *Über die Enden diskreter Räume und Gruppen*, Comment. Math. Helv. **17** (1945), 1–38.
- [7] M.H. Freedman and F. Quinn, *Topology of 4-manifolds*, Princeton Math. Series **39**, Princeton University Press 1990.
- [8] M. Iori and R. Piergallini, *4-manifolds as covers of  $S^4$  branched over non-singular surfaces*, Geometry & Topology **6** (2002), 393–401.
- [9] R. Kirby, *Open problems in low-dimensional topology*, manuscript 1995, math.berkeley.edu/~kirby
- [10] F. Laudenbach and V. Poenaru, *A note on 4-dimensional handlebodies*, Bull. Soc. Math. France **100** (1972), 337–344.
- [11] J.M. Montesinos, *4-manifolds, 3-fold covering spaces and ribbons*, Trans. Amer. Math. Soc. **245** (1978), 453–467.
- [12] J.M. Montesinos, *A note on moves and irregular coverings of  $S^4$* , Contemp. Math. **44** (1985), 345–349.
- [13] J.M. Montesinos, *Representing open 3-manifolds as 3-fold branched coverings*, Rev. Mat. Complut. **15** (2002), 533–542.
- [14] J.M. Montesinos, *Branched coverings after Fox*, Bol. Soc. Mat. Mexicana **11** (2005), 19–64.
- [15] R.P. Osborne, *Embedding Cantor sets in a manifold*, Mich. Math. J. **13** (1966), 57–63.
- [16] R. Piergallini, *Covering Moves*, Trans Amer. Math. Soc. **325** (1991), 903–920.
- [17] R. Piergallini, *Four-manifolds as 4-fold branched covers of  $S^4$* , Topology **34** (1995), 497–508.
- [18] G. Pollini, *Topological 4-manifolds as branched covers*, PhD thesis, Università di Roma “La Sapienza”, 2007.
- [19] C.P. Rourke and B.J. Sanderson, *Introduction to piecewise-linear topology*, Ergebnisse der Mathematik und ihrer Grenzgebiete **69**, Springer-Verlag 1972.